



Herpes simplex virus type 1 encephalitis is associated with elevated levels of F₂-isoprostanes and F₄-neuroprostanes

Dejan Milatovic,¹ Yueli Zhang,¹ Sandra J Olson,¹ Kathleen S Montine,¹ L Jackson Roberts II,^{2,3} Jason D Morrow,^{2,3} Thomas J Montine,^{1,3} Terence S Dermody,^{4,5,6} and Tibor Valyi-Nagy^{1,6}

¹Department of Pathology; ²Department of Medicine; ³Department of Pharmacology; ⁴Department of Microbiology and Immunology; ⁵Department of Pediatrics; and the ⁶Elizabeth B. Lamb Center for Pediatric Research, Vanderbilt University School of Medicine, Nashville, Tennessee, USA

To better understand the pathogenesis of herpes simplex virus type 1 (HSV-1) infections of the nervous system, concentrations of F₄-neuroprostanes (F₄-NP) and F₂-isoprostanes (F₂-IP) in the murine brain were determined following intracerebral inoculation of HSV-1 or normal saline. F₄-NP are highly selective, quantitative markers of neuronal oxidative damage, while F₂-IP are markers of oxidative damage to brain tissue not limited to a certain cell type. In contrast to saline-treated control animals, HSV-1-infected animals developed encephalitic symptoms associated with severe inflammation, widespread HSV-1 protein expression, and significantly elevated F₄-NP and F₂-IP levels in the brain. Survivors of acute HSV-1 infection showed no encephalitic symptoms 2 to 3 weeks following virus inoculation. Brain tissue derived from mice euthanized 2 month after virus inoculation demonstrated expression of HSV-1 latency-associated transcripts without detectable HSV-1 protein expression. However, brain tissue from these animals showed focal chronic inflammation, moderately elevated F₂-IP levels, and normal levels of F₄-NP. These observations provide novel biochemical evidence that oxidant tissue injury is a mechanism underlying neuronal damage during acute HSV-1 encephalitis and suggest that oxidative damage to tissue may continue in the mammalian brain until at least several weeks after recovery from the symptomatic phase of HSV-1 infection. *Journal of NeuroVirology* (2002) **8**, 295–305.

Keywords: HSV-1 encephalitis; oxidative stress; isoprostanes; neuroprostanes; neuroinflammation; neurodegeneration; apoptosis

Introduction

Herpes simplex virus type 1 (HSV-1) is an important pathogen infecting the vast majority of humans

worldwide (reviewed in Whitley, 2001). Following primary infection of mucocutaneous sites, HSV-1 travels by axonal routes to the nervous system where it can cause encephalitis, or much more commonly establishes a latent infection in neurons (reviewed in Skoldenberg, 1996; Steiner, 1996; Whitley, 2001). HSV-1 latent infection of the peripheral nervous system (PNS) affects a majority of humans by adulthood (Liedke *et al*, 1993; Baringer and Pisani, 1994; Schultz *et al*, 1998). Reactivation of HSV-1 from neurons can lead to recurrent mucocutaneous disease and occasionally encephalitis (reviewed in Skoldenberg, 1996; Whitley, 2001). As a consequence of primary infection, HSV-1 genomic DNA is also present in the central nervous system (CNS) of a significant

Address correspondence to Tibor Valyi-Nagy at his current address: University of Illinois at Chicago, College of Medicine, Department of Pathology (MC 847), Room 446, 1819 W. Polk Street, Chicago, IL 60612-7335, USA. E-mail: tiborv@uic.edu

This work was supported by Public Health Service awards R01 AG16835 from the National Institute of Aging (TJM), and R01 AI38296 from the National Institute of Allergy and Infectious Diseases (TSD), the Alzheimer's Disease and Association (TJM), and the Elizabeth B Lamb Center for Pediatric Research (TSD and TVN).

Received 5 February 2002; revised 18 March 2002; accepted 27 March 2002.

proportion of the adult human population (Fraser *et al*, 1981; Liedke *et al*, 1993; Baringer and Pisani, 1994; Itzhaki *et al*, 1997). However, based primarily on observations made using animal models of HSV-1 pathogenesis, it is thought that viral reactivation does not occur in the CNS (Cabrera *et al*, 1980; Steiner *et al*, 1994). Although encephalitis develops in only a small minority of primary and reactivation HSV-1 infections, HSV-1 is the most common cause of sporadic encephalitis in the United States. The molecular mechanisms by which HSV-1 damages the nervous system are not well understood.

Injury mediated by free radicals, most notably by reactive oxygen species (ROS) and reactive nitrogen species (RNS), is an important common pathway of tissue damage for a wide variety of acute and chronic pathologic processes (reviewed in Halliwell and Gutteridge, 1999). ROS and RNS induced by viruses and reactive inflammatory responses can serve as messenger molecules that modulate cellular activation state and influence viral growth. Tissue damage mediated by ROS and RNS is increasingly recognized as an important mechanism by which viral infections can damage the host (reviewed in Schwartz, 1996; Peterhans, 1997a, 1997b; Akaike *et al*, 1998). Free radical-mediated injury also is a mechanism by which viral infections can damage the nervous system. Neural damage induced by free radicals has been reported as a consequence of human immunodeficiency virus infection of humans (Boven *et al*, 1999), reovirus infection of mice (Valyi-Nagy *et al*, 1999), and murine leukemia virus infection of rats (Wilt *et al*, 2000).

ROS and RNS also play important roles in HSV-1 pathogenesis. HSV-1 infection has been found to cause free radical-mediated injury in cultured cells (Palu *et al*, 1994) and in the murine lung (Adler *et al*, 1997). HSV-1 infection induces nitric oxide (NO) production in differentiated monocytes and macrophages in culture (Lopez-Guerrero and Alonso, 1997; Fujioka *et al*, 2000) and in the nervous system of rats (Fujii *et al*, 1999). Infection of ocular, lung, and nervous system tissues of mice is associated with the expression of the inducible form of NO synthetase (iNOS) and the release of cytokines including tumor necrosis factor (TNF)- α from inflammatory cells (Koprowski *et al*, 1993; Adler *et al*, 1997; Shimeld *et al*, 1997; Daheshia *et al*, 1998; Meyding-Lamade *et al*, 1998; Fujii *et al*, 1999; Nucci *et al*, 2000). NO and TNF- α have anti-HSV-1 activity and can either generate potent oxidizing byproducts directly (NO) or induce free radical-mediated injury indirectly through membrane signaling (TNF- α) (Matthews *et al*, 1987; Feduchi *et al*, 1989; Rossol-Voth *et al*, 1991; Croen, 1993; Karupiah *et al*, 1993; Sanchez-Alcazar *et al*, 2000). HSV-1 infections are associated with increased activity of antioxidant enzymes and depletion of the antioxidant glutathione, providing further evidence that HSV-1 infection is associated with oxidative stress (MacLean *et al*, 1998; Nucci *et al*, 2000). In

concordance with this idea, therapy with the antioxidant glutathione reduces the severity of HSV-1 induced eye disease in mice (Nucci *et al*, 2000).

We have studied experimental HSV-1 infections of mice to define the role of free radical-mediated tissue injury in HSV-1 neuropathogenesis. As in humans, infection of peripheral tissues in mice leads to viral replication in epithelia, spread of virus by neural pathways from the PNS to the CNS, and occasional encephalitis depending on viral strain and dose (reviewed in Wagner and Bloom, 1997; Roizman and Knipe, 2001). Direct intracerebral (i.c.) inoculation of HSV-1 also leads to encephalitis. Animals surviving acute HSV-1 infection do not demonstrate signs or symptoms of encephalitis, and infectious virus is not detectable in the nervous system. However, HSV-1 establishes latency in these animals in neurons of the PNS and HSV-1 DNA remains detectable in the CNS (Cabrera *et al*, 1980; Steiner *et al*, 1994). Latent HSV-1 infections of neurons in mice show features similar to those seen in humans: viral genomic DNA is present in neuronal nuclei, abundant viral gene expression is limited to the latency-associated transcripts (LATs), and viral proteins or infectious virus are not detectable (reviewed in Wagner and Bloom, 1997; Roizman and Knipe, 2001). In addition, HSV-1 latency in the nervous system of mice is associated with persistent, chronic inflammation, persistently elevated levels of cytokines, and iNOS activity (Koprowski *et al*, 1993; Cantin *et al*, 1995; Meyding-Lamade *et al*, 1998; Chen *et al*, 2000; Valyi-Nagy *et al*, 2000).

Previous studies in our laboratory using a murine HSV-1 neuropathogenesis model and immunohistochemical detection methods of oxidant tissue injury have provided evidence that free radical-mediated tissue damage is a mechanism by which both acute and latent HSV-1 infections can cause damage in the nervous system (Valyi-Nagy *et al*, 2000). In the present study we used quantitative biochemical detection methods of oxidant tissue injury to test the hypothesis that oxidant brain injury occurs during HSV-1 infection of the murine nervous system. We have shown previously that free radical damage to the brain can be sensitively and accurately quantified by measuring chemically stable oxidative damage products of arachidonic acid (AA) and docosahexaenoic acid (DHA), F₂-isoprostanes (F₂-IP) and F₄-neuroprostanes (F₄-NP), respectively (Morrow and Roberts, 1997; Roberts *et al*, 1998). AA is relatively evenly distributed in the brain with similar concentrations in gray and white matter and within glia and neurons. Thus, measurement of F₂-IP is a reflection of oxidative damage to the entire brain, a limitation shared by all other quantitative methods of determining oxidative damage to the brain, except F₄-NP. Unlike AA, DHA is highly concentrated in neuronal membranes to the exclusion of other cell types. Moreover, we have shown that F₄-NP are by far the most abundant products of this pathway in

the brain (Roberts *et al*, 1998). Thus, measurement of F₄-NP provides a highly selective, quantitative window into neuronal oxidative damage.

We found that levels of F₂-IP and F₄-NP are markedly elevated during acute HSV-1 encephalitis and that levels of F₂-IP remain moderately elevated until several weeks after recovery from the acute, symptomatic phase of infection. These observations provide novel biochemical evidence that oxidant tissue injury is a mechanism underlying neuronal damage during acute HSV-1 encephalitis and suggest that oxidative damage to tissue may continue in the mammalian brain following recovery from the symptomatic phase of HSV-1 infection.

Results

HSV-1 infection of mice

To better understand the pathogenesis of oxidant neural injury during HSV-1 encephalitis, 4- to 6-week-old female BALB/c mice were inoculated i.c. in the left cerebral hemisphere with 1×10^3 or 1×10^5 plaque-forming units (PFU) of HSV-1 strain 17⁺ or normal saline (mock-infection). The i.c. route of inoculation was chosen to provide synchronous, widespread CNS infection. Saline-inoculated animals demonstrated no disease symptoms following recovery from anesthesia. All HSV-1-infected animals developed encephalitic symptoms during the first two weeks following HSV-1-inoculation, irrespective of viral dose. These symptoms included agitation, hunched posture, ruffled fur, lethargy, circling behavior, and paralysis. None of the mice inoculated with 1×10^5 PFU survived beyond 2 weeks following infection; however, occasional mice survived HSV-1 infection following inoculation with 1×10^3 PFU. These surviving animals demonstrated no encephalitic symptoms beyond 2–3 weeks postinoculation and showed normal development.

Acute HSV-1 infection is associated with severe inflammation, widespread HSV-1 protein expression, and significantly elevated F₂-IP and F₄-NP levels in the murine brain

Hematoxylin and eosin-stained sections of tissues derived from the right brain half of mice euthanized 5 to 8 days after i.c. HSV-1 inoculation invariably demonstrated evidence of meningoencephalitis. Histologic sections showed inflammatory cell infiltrates in the brain parenchyma and meninges. Inflammatory changes primarily involved the brain stem, the diencephalon, the cerebral cortex, and the olfactory bulb. Perivascular and parenchymal inflammatory infiltrates consisted mostly of mononuclear inflammatory cells, but occasional necrotizing foci associated with infiltrates rich in neutrophils also were observed (Figure 1A, Table 1). Inflamed foci demonstrated widespread expression of HSV-1 proteins (Figure 1E, Table 1) and LAT (Figure 2A, Table 1). HSV-1 immunoreactivity and LAT expression pri-

marily involved cells with the morphologic features of neurons. Sections derived from the right brain half of saline-inoculated animals euthanized 5 days following inoculation demonstrated normal histology and no evidence of HSV-1 protein or LAT expression (Figures 1B, 1F, 2B, Table 1).

Apoptotic cell death in the brain of HSV-1-infected and mock-infected mice was determined by detection of apoptosis-associated DNA strand breaks in tissue sections using a terminal deoxynucleotidyl transferase (TdT)-mediated DUTP-digoxigenin nick end-labeling (TUNEL) assay. TUNEL analysis of brain tissues derived from mice euthanized 5 to 8 days after HSV-1 inoculation demonstrated numerous TUNEL-positive cells that were present primarily in inflamed brain areas (Figure 3A). Morphologically, TUNEL-positive cells consisted of both neurons and nonneuronal cells. CNS tissues derived from uninfected mice and from mice euthanized five days following mock-infection with saline contained no TUNEL-positive cells (Figure 3B). These findings suggest that acute HSV-1 encephalitis is associated with apoptosis, in agreement with previous reports (Geiger *et al*, 1995, 1997; Pelosi *et al*, 1998; Valyi-Nagy *et al*, 2000).

To determine whether oxidant injury occurs during acute HSV-1 infection of the murine CNS, concentrations of F₂-IP and F₄-NP in the brain of mice were determined following i.c. inoculation of HSV-1 or normal saline (Figures 4, 5, Table 1). F₂-IP levels did not increase in the left, inoculated brain half of saline-treated animals euthanized 5 days after inoculation relative to F₂-IP levels detected in the brain of untreated mice (Figure 4, Table 1). This finding indicates that the minor trauma associated with inoculation does not alter F₂-IP levels. Brain tissues of mice euthanized 5 to 8 days after HSV-1 inoculation demonstrated significantly elevated F₂-IP levels relative to saline-inoculated ($P < 0.01$) and untreated controls ($P < 0.01$) (Figure 4, Table 1). Brain tissues of mice euthanized on days 5 to 8 after inoculation with HSV-1 also showed significantly elevated F₄-NP levels relative to saline-inoculated controls ($P < 0.02$) (Figure 5, Table 1). These findings demonstrate that i.c. inoculation of HSV-1 into mice leads to generalized encephalitis associated with oxidant brain injury that involves neurons.

Evidence of focal inflammation and moderately elevated F₂-IP levels in the murine brain several weeks after recovery from symptomatic HSV-1 encephalitis

To determine whether inflammation, evidence of HSV-1 infection, or oxidant tissue injury are detectable in the CNS of mice that survive the acute, symptomatic phase of HSV-1 infection, groups of mice were euthanized 60 days after i.c. inoculation with either HSV-1 or normal saline. Hematoxylin and eosin-stained sections of tissues derived from the right brain half of five of six mice euthanized 60 days after i.c. HSV-1 inoculation demonstrated




Figure 1 Inflammation and HSV-1 protein expression in the right cerebral hemisphere of BALB/c mice following i.c. inoculation of either HSV-1 or normal saline (mock-infection) into the left cerebral hemisphere. Hematoxylin and eosin-stained sections (A) 5 days following inoculation of 10^5 PFU of HSV-1. (B) 5 days following mock-infection. (C) 60 days following inoculation of 10^3 PFU of HSV-1. (D) 60 days following mock-infection. Detection of HSV-1 protein expression using a polyclonal anti-HSV-1 antiserum. (E) 5 days following inoculation of 10^5 PFU of HSV-1. (F) 5 days following mock-infection. (G) 60 days following inoculation of 10^3 PFU of HSV-1. (H) 60 days following mock-infection. (magnification 400 \times).

Table 1 HSV-1 protein and LAT expression and inflammation in the right brain half and F₂-isoprostane and F₄-neuroprostane levels in the left brain half of adult BALB/c mice 5 to 8 days and 60 days following intracerebral (i.c.) inoculation of HSV-1 strain 17⁺ or normal saline

Inoculum (i.c.)	Time of euthanasia (days after i.c. inoculation)	No. of animals pos/No. of animals tested for:			Brain concentration of:	
		HSV-1 protein expression ¹	HSV-1 LAT expression ²	Inflammation	F ₂ -isoprostanes (ng/g) ± SD	F ₄ -neuroprostanes (ng/g) ± SD
None	NA ³	0/6	0/6	0/6	2.77 ± 0.28 n = 9	ND ⁴
Saline	5	0/3	0/3	0/3	2.70 ± 0.15 n = 6	60.2 ± 8.1 n = 8
HSV-1 (10 ³ –10 ⁵ PFU)	5–8	10/10	5/5	10/10	5.82 ± 1.02* n = 8	132.8 ± 16.0* n = 5
Saline	60	0/5	0/5	0/5	2.52 ± 0.29 n = 4	57.0 ± 14.7 n = 4
HSV-1 (10 ³ PFU)	60	0/6	5/6	5/6	3.71 ± 0.44* n = 6	67.0 ± 22.3 n = 6

¹As detected by immunohistochemistry.²As detected by *in situ* hybridization.³NA = not applicable.⁴ND = not done.**P* < 0.05 compared to saline.

evidence of focal chronic inflammation in brain tissues (Table 1). Inflammatory changes were less extensive than those detected in tissues derived from animals euthanized 5 to 8 days after HSV-1 inoculation but showed similar distribution in the nervous system. Inflammatory cell infiltrates were composed of mononuclear cells and involved both brain parenchyma and meninges (Figure 1C). HSV-1 protein expression was not detected by immunohistochemistry in sections of brain tissues derived from mice euthanized 60 days after inoculation (Figure 1G, Table 1). However, HSV-1 LAT expression was detected by *in situ* hybridization in the brain of five of six mice examined (Figure 2C, Table 1). LAT expression primarily involved cells with the morphologic features of neurons. Because LAT expression occurs during both productive and latent infection, and HSV-1 protein expression occurs only during productive infection (Wagner and Bloom, 1997; Roizman and Knipe, 2001), these findings indicate that productive HSV-1 infection had subsided in the brain of mice by 60 days following virus inoculation. Sections derived from brain tissue of control animals euthanized 60 days following saline inoculation demonstrated normal histology and no evidence of HSV-1 protein or LAT expression (Figures 1D, 1H, 2D, Table 1).

TUNEL analysis of brain tissues derived from mice euthanized sixty days after HSV-1 inoculation showed occasional TUNEL-positive cells that were present primarily in inflamed brain areas (Figure 3C). Morphologically, TUNEL-positive cells consisted of both neurons and nonneuronal cells. CNS tissues derived from mice euthanized 60 days following mock-infection with saline contained no TUNEL-positive cells (Figure 3D). These findings suggest that HSV-1 infection is associated with apoptosis in the murine

brain several weeks after recovery from the symptomatic phase of acute encephalitis.

Brain tissue of mice euthanized 60 days after HSV-1 inoculation demonstrated significantly elevated F₂-IP levels relative to mock-infected (*P* = 0.04) and untreated controls (*P* = 0.04) (Figure 4). However, F₂-IP levels in brain tissue of mice 60 days after HSV-1 inoculation were lower than those associated with acute HSV-1 infection. Brain tissue of mice euthanized on day 60 postinoculation with HSV-1 demonstrated F₄-NP levels similar to saline-inoculated controls (Figure 5, Table 1). These findings suggest that oxidative damage to tissue may continue in the mammalian brain until at least several weeks after recovery from the symptomatic phase of HSV-1 encephalitis and that this injury primarily involves nonneuronal cells.

Discussion

In this report, we show that levels of F₂-IP and F₄-NP, which are accurate *in vivo* biomarkers of oxidant injury, are significantly elevated during acute HSV-1 encephalitis. In addition, we show that levels of F₂-IP remain moderately elevated in the murine brain until at least several weeks after recovery from the symptomatic phase of HSV-1 infection. Following acute infection, F₄-NP levels normalize, indicating that nonneuronal cells are the primary targets of oxidative neural injury in mice that survive acute HSV-1 encephalitis.

The demonstration by biochemical methods that HSV-1 infection is associated with oxidant brain injury confirms and extends our previous studies using immunohistochemical methods (Valyi-Nagy *et al*, 2000). As F₄-NP are highly selective, quantitative markers of neuronal oxidative damage, while F₂-IP







Figure 2 HSV-1 LAT expression in the right side of the brain stem of BALB/c mice following i.c. inoculation of either HSV-1 or normal saline (mock-infection) into the left cerebral hemisphere. HSV-1 LAT expression (A) 5 days following inoculation of 10^5 PFU of HSV-1. (B) 5 days following mock-infection. (C) 60 days following mock-infection. (D) 60 days following HSV-1. (Magnification 200 \times .)




Figure 3 TUNEL-positive cells in the right cerebral hemisphere of BALB/c mice following i.c. inoculation of either HSV-1 or normal saline (mock-infection) into the left cerebral hemisphere. Sections were analyzed by TUNEL assay to detect apoptosis-associated DNA strand breaks (A) 5 days following inoculation of 10^5 PFU of HSV-1. (B) 5 days following mock-infection. (C) 60 days following inoculation of 10^3 PFU of HSV-1. (D) 60 days following mock-infection. (Magnification 400 \times .)




Figure 4 F₂-isoprostane (F₂-IP) concentrations in the left brain half of BALB/c mice following i.c. inoculation of either HSV-1 or normal saline (mock-infection) into the left cerebral hemisphere. Untreated: 4- to 6-week-old mice without treatment. Mock 5d: mice euthanized 5 days following mock-infection. HSV 5-8 d: six mice euthanized 5 days following inoculation of 10^5 PFU of HSV-1 and two mice following inoculation of 10^3 PFU of HSV-1. Mock 2 mo: mice euthanized 60 days after mock-infection. HSV 2 mo: mice euthanized 60 days following inoculation of 10^3 PFU of HSV-1. Numbers above the bars refer to the number of animals in the indicated group. *indicates $P < 0.05$ compared to mock-infection.




Figure 5 F₄-neuroprostane (F₄-NP) concentrations in the left brain half of BALB/c mice following i.c. inoculation of either HSV-1 or normal saline (mock-infection) into the left cerebral hemisphere. Mock acute: mice euthanized 5 days following mock-infection. HSV acute: three mice euthanized 5 days and two mice euthanized 8 days following inoculation of 10^3 PFU of HSV-1. Mock 2 mo: mice euthanized 60 days following mock-infection. HSV 2 mo: mice euthanized 60 days following inoculation of 10^3 PFU of HSV-1. Numbers above the bars refer to the number of animals in the indicated group. *indicates $P < 0.05$ compared to mock-infection.

are markers of oxidative damage to brain tissue not limited to a certain cell type, our findings indicate that oxidant injury is a mechanism of neuronal damage during acute HSV-1 encephalitis. These results do not identify specific viral or host factors responsible for oxidant brain injury during HSV-1 encephalitis. However, as HSV-1 infection has been found to cause free radical-mediated injury in cultured cells (Palu *et al*, 1994), it appears possible that both viral and inflammatory mediators are responsible for the observed effects.

Steady-state levels of free radical-mediated tissue damage represent a balance between rates of damage caused by pro-oxidant stimuli and rates of antioxidant and tissue repair mechanisms that decrease ROS and RNS levels and remove oxidatively damaged molecules (Halliwell and Gutteridge, 1999). Observations from our laboratory indicate that the turnover of F₂-IP and F₄-NP in the mammalian brain is quite rapid. In the rat hippocampus, markedly elevated F₂-IP levels associated with kainate-induced seizures return to baseline levels by 4 days after kainate injection (Patel *et al*, 2001). Furthermore, we have found that increased F₂-IP and F₄-NP levels induced by intracerebroventricular inoculation of bacterial lipopolysaccharide return to normal by 2 to 9 days after inoculation (Milatovic *et al*, unpublished observation). These findings strongly suggest that the detection of elevated F₂-IP in brain tissue of mice several weeks after recovery from symptomatic HSV-1 encephalitis is not due to residual damage from the acute phase of infection. Furthermore, the detected divergence of F₂-IP and F₄-NP levels 2 months after virus inoculation suggests that oxidant brain injury following resolution of HSV-1 encephalitis primarily affects nonneuronal cells. As we found no evidence of HSV-1 protein expression in the brain of mice 2 months after virus inoculation, it appears likely that inflammatory mediators rather than direct viral toxicity are primarily responsible for the observed effects.

HSV-1 encephalitis in the murine brain is associated with apoptosis of neurons and nonneuronal cells (Geiger *et al*, 1995, 1997; Pelosi *et al*, 1998; Valyi-Nagy *et al*, 2000). Cells undergoing apoptosis in the setting of HSV-1 encephalitis colocalize with sites of virus infection and oxidative damage in the brain (Valyi-Nagy *et al*, 2000). Findings reported here indicate that elevated F₂-IP and F₄-NP levels during acute HSV-1 infection are associated with increased apoptosis in the CNS. Furthermore, moderately elevated F₂-IP levels 2 months after HSV-1 inoculation are associated with moderately increased numbers of apoptotic cells in the murine brain. As oxidative damage can lead to cell death by apoptosis (reviewed by Halliwell and Gutteridge, 1999), these findings raise the possibility that oxidative damage is a determinant of cell death during HSV-1 encephalitis. In addition, these findings suggest that HSV-1 infection-induced oxidative damage may lead to apoptosis in

the mammalian brain several weeks after recovery from the symptomatic phase of acute encephalitis.

Human survivors of HSV-1 encephalitis continue to harbor HSV-1 DNA in the brain and exhibit evidence of chronic inflammation in the absence of virus replication for months to years (Ellison and Love, 1996). It is not known whether carriage of HSV-1 DNA is associated with brain injury in survivors of encephalitis or in those much more numerous individuals also harboring HSV-1 DNA in their brain as a consequence of less severe primary infections. Because a significant proportion of the adult human population contains HSV-1 DNA in the brain, the pathological significance of this infection is important to understand. The presence of HSV-1 DNA in human brain in the context of the apolipoprotein E epsilon 4 allele has been proposed to represent a strong risk factor for the development of the most common human neurodegenerative disease, Alzheimer's disease (Itzhaki *et al*, 1997; Dobson and Itzhaki, 1999). Although an association between HSV-1 infection and Alzheimer's disease remains to be proven, our findings indicate that oxidative destruction of tissue may continue in the mammalian brain until at least several weeks after recovery from the symptomatic phase of HSV-1 infection.

Findings reported here represent to our knowledge the first quantitative biochemical demonstration of free radical-mediated tissue injury induced by a virus in the nervous system. The role of viral and host factors in the pathogenesis of virus-induced oxidant tissue injury in the nervous system is almost completely unknown. Demonstration of the feasibility of quantitative biochemical detection of free radical-mediated tissue injury induced by HSV-1 in the murine nervous system provides a new experimental system for dissection of mechanisms underlying oxidant tissue injury associated with neurotropic viral infections.

Materials and methods

Virus inoculation and tissue collection

4- to 6-week-old female BALB/c mice (Harlan, Indianapolis, IN) were inoculated by the i.c. route in the left cerebral hemisphere with 1×10^3 or 1×10^5 PFU of HSV-1 strain 17⁺ or sterile normal saline (mock-infection) under anesthesia (Valyi-Nagy *et al*, 1994). The inoculum volume was 5 μ L. Mice were observed three times daily for signs and symptoms of disease, and those demonstrating advanced encephalitis (lethargy, paralysis) were euthanized. Survivors of acute HSV-1 infection were euthanized 2 months after inoculation. Groups of saline-inoculated mice were euthanized 5 days and 2 months after mock-infection.

Brain tissue of euthanized mice was removed and divided into right and left halves in the midsagittal plane. Tissues derived from both sides contained

cerebrum, olfactory bulb, brain stem, and cerebellum. The left half was flash frozen in liquid nitrogen and then maintained at -80°C ; the right half was fixed in 10% buffered formaldehyde. Frozen tissues corresponding to the inoculated left brain halves were used to measure $\text{F}_2\text{-IP}$ and $\text{F}_4\text{-NP}$ levels. Formaldehyde-fixed tissues were paraffin-embedded and sectioned. Sections were stained with hematoxylin and eosin, incubated with a polyclonal anti-HSV-1 serum, or analyzed by *in situ* hybridization for HSV-1 LAT expression.

Immunohistochemistry

Tissue sections for immunohistochemistry were deparaffinized with xylene and rehydrated through a series of graded ethanols. Endogenous peroxidase was quenched in a 0.3% H_2O_2 -methanol bath followed by several washes with phosphate-buffered saline (PBS). Prior to staining, binding of secondary antibodies and conjugates was blocked by appropriate reagents provided by the manufacturer. HSV-1 proteins were detected using a 1:1000 dilution of a polyclonal anti-HSV-1 antiserum raised in a rabbit (Dako, Carpinteria, CA). Tissue sections were incubated with primary antibody at 43°C for 32 min prior to addition of biotinylated anti-rabbit-immunoglobulin secondary antibody, avidin-horse-radish-peroxidase, and 3,3'-diaminobenzidine

tetrahydrochloride (0.04%) in 0.05 M Tris-HCl (pH 7.4) and 0.025% H_2O_2 as a chromogen (Ventana Medical Systems, Tucson, AZ).

In situ hybridization *In situ* hybridization was performed using a nick-translated ^{35}S -labeled DNA probe specific for HSV-1 LAT (Bst2-Bst2 fragment) (Valyi-Nagy *et al*, 1992) using previously described techniques (Valyi-Nagy *et al*, 1991). The probe used in these experiments had a specific activity of $>10^8$ cpm per microgram.

Detection of apoptosis Apoptotic cell death in the brain of HSV-1-infected and mock-infected mice was determined by detection of apoptosis-associated DNA strand breaks in tissue sections using a TUNEL assay (ApopTag Plus; Onco, Gaithersburg, MD).

Quantification of F_2 -isoprostanes ($\text{F}_2\text{-IP}$) and F_4 -neuroprostanes ($\text{F}_4\text{-NP}$) A stable isotope dilution method followed by gas chromatography and negative ion chemical ionization mass spectrometry was used to determine $\text{F}_2\text{-IP}$ and $\text{F}_4\text{-NP}$ concentrations in the frozen left brain half of euthanized mice according to previously described methods (Morrow and Roberts, 1997; Roberts *et al*, 1998).

Data analysis Tissue sections processed for immunohistochemistry and *in situ* hybridization were reviewed by an observer blinded to the nature of the infection (HSV-1 or mock). Results were analyzed using the one-tailed *t*-test.

References

- Adler H, Beland JL, Del-Pan NC, Kobzik L, Brewer JP, Martin TR, Rimm IJ (1997). Suppression of herpes simplex virus type 1 (HSV-1)-induced pneumonia in mice by inhibition of inducible nitric oxide synthase (iNOS, NOS2). *J Exp Med* **185**: 1533–1540.
- Akaike T, Suga M, Maeda H (1998). Free radicals in viral pathogenesis: Molecular mechanisms involving superoxide and NO. *Proc Soc Ex Biol Med* **217**: 64–73.
- Baringer JR, Pisani P (1994). Herpes simplex virus genomes in human nervous system tissues analyzed by polymerase chain-reaction. *Ann Neurol* **36**: 823–829.
- Boven LA, Gomes L, Hery C, Gray F, Verhoef J, Portegies P, Tardieu M, Nottet HS (1999). Increased peroxynitrite activity in AIDS dementia complex: Implications for neuropathogenesis of HIV-1 infection. *J Immunol* **162**: 4319–4327.
- Cabrera CV, Wohlenberg C, Openshaw H, Rey Mendez M, Puga A, Notkins AL (1980). Herpes simplex virus DNA sequences in the CNS of latently infected mice. *Nature* **288**: 288–290.
- Cantin EM, Hinton DR, Chen J, Openshaw H (1995). Gamma interferon expression during acute and latent nervous system infection by herpes simplex virus type 1. *J Virol* **69**: 4898–4905.
- Chen SH, Garber DA, Schaffer PA, Knipe DM, Coen DM (2000). Persistent elevated expression of cytokine transcripts in ganglia latently infected with herpes simplex virus in the absence of ganglionic replication or reactivation. *Virology* **278**: 207–216.
- Croen D (1993). Evidence for antiviral effect of nitric oxide. Inhibition of herpes simplex virus type 1 replication. *J Clin Invest* **91**: 2446–2452.
- Daheshia M, Kanangat S, Rouse BT (1998). Production of key molecules by ocular neutrophils early after herpetic infection of the cornea. *Exp Eye Res* **67**: 619–624.
- Dobson CB, Itzhaki RF (1999). Herpes simplex virus type 1 and Alzheimer's disease. *Neurobiol Aging* **20**: 457–465.
- Ellison D, Love S (1996). Acute viral infections. In *Neuropathology*. Ellison D, Love S (eds). Mosby (Harcourt Publishers Limited): London, pp 12.1–12.19.
- Feduchi E, Alonso MA, Carrasco L (1989). Human gamma interferon and tumor necrosis factor exert a synergistic blockade on the replication of herpes simplex virus. *J Virol* **63**: 1354–1359.
- Fraser NW, Lawrence WC, Wroblewska Z, Gilden DH, Koprowski H (1981). Herpes simplex type 1 DNA in human brain tissue. *Proc Natl Acad Sci USA* **78**: 6461–6465.
- Fujii S, Akaike T, Maeda H (1999). Role of nitric oxide in the pathogenesis of herpes simplex virus encephalitis in rats. *Virology* **256**: 203–212.
- Fujioka N, Ohashi K, Ikeda M, Kurimoto M (2000). Autocrine interferon-beta stimulation augments nitric oxide production by mouse macrophage J774A.1 cells infected with herpes simplex virus type 1. *Microbiol Immunol* **44**: 283–287.
- Geiger DK, Gurushanthaiah D, Howes EL, Lewandowski GA, Reed JC, Bloom FE, Sarvetnick NE (1995).

- Cytokine-mediated survival from lethal herpes simplex virus infection: Role of programmed neuronal death. *Proc Natl Acad Sci USA* **92**: 3411–3415.
- Geiger DK, Nash TC, Sawyer S, Krahl T, Patstone G, Reed JC, Krajewski S, Dalton D, Buchmeier MJ, Sarvetnick NE (1997). Interferon- γ protects against herpes simplex virus type 1-mediated neuronal death. *Virology* **238**: 189–197.
- Halliwell B, Gutteridge JM (1999). *Free radicals in biology and medicine*. 3rd ed. Oxford Univ Press: Oxford, UK.
- Itzhaki RF, Lin WR, Shang D, Wilcock GK, Faragher B, Jamieson GA (1997). Herpes simplex virus type 1 in brain and risk of Alzheimer's disease. *Lancet* **349**: 241–244.
- Karupiah G, Xie QW, Buller RM, Nathan C, Duarte C, MacMicking JD (1993). Inhibition of viral replication by interferon-gamma-induced nitric oxide synthase. *Science* **261**: 1445–1448.
- Koprowski H, Zheng YM, Heber-Katz E, Fraser N, Rorke L, Fu ZF, Hanlon C, Dietzschold B (1993). *In vivo* expression of inducible nitric oxide synthetase in experimentally induced neurologic diseases. *Proc Natl Acad Sci USA* **90**: 3024–3027.
- Liedke W, Opalka B, Zimmermann CW, Lignitz E (1993). Age distribution of latent herpes simplex virus 1 and varicella-zoster virus genome in human nervous tissue. *J Neurol Sci* **116**: 6–11.
- Lopez-Guerrero JA, Alonso MA (1997). Nitric oxide production induced by herpes simplex virus type 1 does not alter the course of the infection in human monocytic cells. *J Gen Virol* **78**: 1977–1980.
- MacLean A, Wei XQ, Huang FP, Al-Alem UA, Chan WL, Liew FY (1998). Mice lacking inducible nitric-oxide synthase are more susceptible to herpes simplex virus infection despite enhanced Th1 cell responses. *J Gen Virol* **79**: 825–830.
- Meyding-Lamade U, Hass J, Lamade W, Stingle K, Kehm R, Fath A, Heinrich K, Storch Hagenlocher B, Wildemann B (1998). Herpes simplex virus encephalitis: Long-term comparative study of viral load and the expression of immunologic nitric-oxide synthetase in mouse brain tissue. *Neurosci Lett* **244**: 9–12.
- Matthews N, Neale ML, Jackson SK, Stark JM (1987). Tumour cell killing by tumour necrosis factor: Inhibition by anaerobic conditions, free radical scavengers and inhibitors of arachidonate metabolism. *Immunology* **62**: 153–155.
- Morrow JD, Roberts LJ (1997). The isoprostanes: Unique bioactive products of lipid peroxidation. *Prog Lipid Res* **36**: 1–21.
- Nucci C, Palamara AT, Ciriolo MR, Nencioni L, Savini P, D'Agostini C, Rotilio G, Cerulli L, Garaci E (2000). Imbalance in corneal redox state during herpes simplex virus 1-induced keratitis in rabbits. Effectiveness of exogenous glutathione supply. *Exp Eye Res* **70**: 215–220.
- Palu G, Biasolo MA, Sartor G, Masotti L, Papini E, Floreani M, Palatini P (1994). Effects of herpes simplex virus type 1 infection on the plasma membrane and related functions of HeLa S3 cells. *J Gen Virol* **75**: 3337–3344.
- Patel M, Liang L, Roberts LJ (2001). Enhanced hippocampal F₂-isoprostane formation following kainate-induced seizures. *J Neurochem* **79**: 1065–1069.
- Pelosi E, Rozenberg F, Coen DM, Tyler KL (1998). A herpes simplex virus DNA polymerase mutation that specifically attenuates neurovirulence in mice. *Virology* **252**: 364–372.
- Peterhans E (1997a). Reactive oxygen species and nitric oxide in viral diseases. *Biol Trace Elem Res* **56**: 107–116.
- Peterhans E (1997b). Oxidants and antioxidants in viral diseases: Disease mechanisms and metabolic regulation. *J Nutri* **127**: 962S–965S.
- Roberts LJ, Montine TJ, Markesbery WR, Tapper AR, Hardy P, Chemtob S, Dettbarn WD, Morrow JD (1998). Formation of isoprostane-like compounds (neuroprostanes) *in vivo* from docosahexaenoic acid. *J Biol Chem* **273**: 13605–13612.
- Roizman B, Knipe DM (2001). Herpes simplex viruses and their replication. In *Fields' Virology*. Knipe DM, Howley PM (eds). Lippincott-Raven: Philadelphia, pp 2399–2460.
- Rossol-Voth R, Rossol S, Schut KH, Corridori S, deCian W, Falke D (1991). *In vivo* protective effect of tumour necrosis factor alpha against experimental infection with herpes simplex virus type 1. *J Gen Virol* **72**: 143–147.
- Sanchez-Alcazar JA, Schneider E, Martinez MA, Carmona P, Hernandez-Munoz I, Siles E, De La Torre P, Ruiz-Cabello J, Garcia I, Solis-Herruzo JA (2000). Tumour necrosis factor-alpha increases the steady-state reduction of cytochrome b of the mitochondrial respiratory chain in metabolically inhibited L929 cells. *J Biol Chem* **275**: 13353–13361.
- Schultz P, Arbusow V, Strupp M, Dieterich M, Rauch E, Brandt T (1998). Highly variable distribution of HSV-1-specific DNA in human geniculate, vestibular and spiral ganglia. *Neurosci Lett* **252**: 139–142.
- Schwarz KB (1996). Oxidative stress during viral infection: A review. *Free Radic Biol Med* **21**: 641–649.
- Shimeld C, Whiteland JL, Williams NA, Easty DL, Hill TJ (1997). Cytokine production in the nervous system of mice during acute and latent infection with herpes simplex virus type 1. *J Gen Virol* **78**: 3317–3325.
- Skoldenberg B (1996). Herpes simplex encephalitis. *Scan J Infect Dis Suppl* **100**: 8–13.
- Steiner I (1996). Human herpesviruses latent infection in the nervous system. *Immunol Rev* **152**: 157–173.
- Steiner I, Modor N, Reibstein I, Spivack JG, Fraser NW (1994). Herpes simplex virus type 1 gene expression and reactivation of latent infection in the central nervous system. *Neuropathol Appl Neurobiol* **20**: 253–260.
- Valyi-Nagy T, Deshmane S, Dillner A, Fraser NW (1991). Induction of cellular transcription factors in trigeminal ganglia of mice by corneal scarification, herpes simplex virus type 1 infection and explantation of trigeminal ganglia. *J Virol* **65**: 4142–4152.
- Valyi-Nagy T, Deshmane SL, Raengsakulrach B, Nicosia M, Gesser RM, Wysocka M, Dillner A, Fraser NW (1992). Herpes simplex virus type 1 mutant in 1814 establishes a unique, slowly progressing infection in SCID mice. *J Virol* **66**: 7336–7345.
- Valyi-Nagy T, Fareed MU, O'Keefe JS, Gesser RM, MacLean AR, Brown SM, Spivack JG, Fraser NW (1994). An HSV-1 strain 17⁺ gamma 34.5 deletion mutant 1716 is avirulent in SCID mice. *J Gen Virol* **75**: 2059–2063.
- Valyi-Nagy T, Olson SJ, Valyi-Nagy K, Montine TJ, Dermody TS (2000). Herpes simplex virus type 1 latency in the murine nervous system is associated with oxidative damage to neurons. *Virology* **278**: 309–321.
- Valyi-Nagy T, Sidell K, Marnett LJ, Dermody TS, Roberts LJ, Morrow JD, Montine TJ (1999). Divergence of brain

- prostaglandin H synthetase activity and oxidative damage in mice with encephalitis. *J Neuropathol Exp Neurol* **58**: 1269–1272.
- Wagner EK, Bloom DC (1997). Experimental investigation of herpes simplex virus latency. *Clin Microbiol Rev* **10**: 419–434.
- Whitley RJ (2001). Herpes simplex viruses. In *Fields' Virology*. Knipe DM, Howley PM (eds). Lippincott-Raven Press: Philadelphia, pp 2461–2510.
- Wilt SG, Dugger NV, Hitt ND, Hoffman PM (2000). Evidence of oxidative damage in a murine leukemia virus-induced neurodegeneration. *J Neurosci Res* **62**: 440–450.